

Geometric phase due to helicity inversion of photons inside an optical fiber composed periodically of left- and right- handed media

Jian-Qi Shen^{1,2 *}

¹ Centre for Optical and Electromagnetic Research, State Key Laboratory of Modern Optical Instrumentation, Zhejiang University, Hangzhou Yuquan 310027, P. R. China

² Zhejiang Institute of Modern Physics and Department of Physics, Zhejiang University, Hangzhou 310027, P. R. China
(February 2, 2008)

In this Letter, it is claimed that, in addition to the Chiao-Wu geometric phase and optical Aharonov-Carmi geometric phase, there exists a new interesting geometric phase caused by the helicity inversion of photons in the optical fiber composed periodically of left- and right- handed (LRH) media. By making use of the Lewis-Riesenfeld invariant theory and the invariant-related unitary transformation formulation, we calculate this geometric phase. It is emphasized that this geometric phase should not be neglected when considering the anomalous refraction on the interfaces between left- and right- handed media.

Keywords: Geometric phase, helicity inversion, left-handed media

More recently, a kind of composite media (the so-called *left-handed media*) having a frequency band where the effective permittivity (ϵ) and the effective permeability (μ) are *simultaneously negative* attracts attention of many researchers in various fields such as materials science, condensed matter physics, optics and electromagnetism [1–5]. In 1964¹, Veselago considered this peculiar medium and showed that it possesses a *negative* index of refraction [6]. It follows from the Maxwell’s equations that in this medium the Poynting vector and wave vector of electromagnetic wave would be antiparallel, *i. e.*, the wave vector $\mathbf{k}$, the electric field $\mathbf{E}$ and the magnetic field $\mathbf{H}$ form a *left-handed* system; thus Veselago referred to such materials as “*left-handed* (LH) ”, and correspondingly, the ordinary medium in which $\mathbf{k}$, $\mathbf{E}$ and $\mathbf{H}$ form a right-handed system may be termed the “*right-handed* (RH) ” medium. Other authors call this class of materials “negative-index media (NIM)”, “double negative media (DNM) [5]” and Veselago’s media. It is readily verified that in such media having both ϵ and μ *negative*, there exist a number of peculiar electromagnetic properties, for instance, many dramatically different propagation characteristics stem from the sign change of the group velocity, including reversal of both the Doppler shift and Cherenkov radiation, anomalous refraction, modified spontaneous emission rates and even reversals of radiation pressure to radiation tension [2,6]. In experiments, this artificial negative electric permittivity media may be obtained by using the *array of long metallic wires* (ALMWs), which simulates the plasma behavior at microwave frequencies, and the artificial negative magnetic permeability media may be built up by using small resonant metallic particles, *e. g.*, the *split ring resonators* (SRRs) with very high magnetic polarizability [3,7–9].

All the above optical and electromagnetic properties and effects of left-handed media were predicted and studied based on the classical Maxwell’s theory. Since left-handed media possess so many peculiar properties and effects, we think it is also physically interesting to consider some certain quantum effects in the left-handed optical fiber. So, in this paper, we deal with a problem of geometric quantum phase of photons inside the left- and right- handed (LRH)-periodical fiber.

Historically, the first physical realization of the Berry’s quantum geometric (topological) phase [10] was proposed by Chiao and Wu [11–14] after Berry suggested the adiabatic quantum theory. By using Berry’s adiabatic phase formula [10], Chiao and Wu investigated the geometric phase of the polarized photon propagating inside a noncoplanarly curved optical fiber [11]. Based on the Lewis-Riesenfeld invariant theory [15–17], we have calculated the non-adiabatic non-cyclic Chiao-Wu geometric phase of photons [18]. Since it is readily verified that the photon propagating inside a curved fiber may be regarded as acted upon by an effective Coriolis force (Lorentz-type force) [18,19], one can predict that there exists an optical Aharonov-Carmi effect [20,21] for moving photons in the curved optical fiber, where the

*E-mail address: jqshen@coer.zju.edu.cn

¹Note that, in the literature, many authors mention the year when Veselago suggested the *left-handed media* by mistake. They claim that Veselago proposed the concept of *left-handed media* in 1968. On the contrary, the true fact is as follows: Veselago’s excellent paper was first published in July, 1964 [Usp. Fiz. Nauk **92**, 517-526 (1964)]. In 1968, this original paper was translated into English by W. H. Furry and published again in the journal of Sov. Phys. Usp. [6].

geometric phase is similar to the Aharonov-Bohm phase.

In the present Letter, we argue that, if, for example, the photons propagating inside a fiber that is composed periodically of left- and right- handed (LRH) media, then a new geometric phase of photons due to helicity inversion will arise. In this LRH-periodical optical fiber, helicity inversion (or the transitions between helicity states) of photons may be easily caused by the interaction of light field with interfaces of LRH materials. So, this geometric phase arises on the interfaces between left- and right- handed media (LRH interfaces), where the anomalous refraction occurs when the incident lightwave travels to the LRH interfaces. we think, in the literature, it gets less attention than it deserves. So, in what follows, we calculate the photon wavefunction and geometric phase in this physical process, and emphasize that we should attach importance to this geometric phase when considering the wave propagation on the LRH interfaces.

Let us consider the applied optical fiber that is fabricated periodically from both left- and right- handed (LRH) media with the optical refractive index being $-n$ and n , respectively. Thus the wave vector of photon moving along the fiber is respectively $-n\frac{\omega}{c}$ in LH section and $n\frac{\omega}{c}$ in RH section, where ω and c respectively denote the frequency and speed in a vacuum of light. For convenience, we assume that the periodical length, b , of LH is equal to that of RH in the fiber. If the eigenvalue of photon helicity is σ in right-handed sections, then, according to the definition of helicity, $h = \frac{\mathbf{k}}{|\mathbf{k}|} \cdot \mathbf{J}$ with $\mathbf{J}$ denoting the total angular momentum of the photon, the eigenvalue of helicity acquires a minus sign in left-handed sections. We assume that at $t = 0$ the light propagates in the right-handed section and the initial eigenvalue of photon helicity is σ . So, in the wave propagation inside the LRH-periodical optical fiber, the helicity eigenvalue of h is then $(-)^m \sigma$ with $m = [\frac{ct}{nb}]$, where $[\]$ represents the integer part of $\frac{ct}{nb}$. This, therefore, means that when $2k(\frac{nb}{c}) < t \leq (2k+1)(\frac{nb}{c})$, $(-)^m = +1$ and when $(2k+1)(\frac{nb}{c}) < t \leq (2k+2)(\frac{nb}{c})$, $(-)^m = -1$, where k is an integer ($k \geq 0$). It follows that the incidence of lightwave in the fiber on the LRH interfaces gives rise to transitions between the helicity states ($|+\rangle$ and $|-\rangle$) of photons. This enables us to construct a time-dependent effective Hamiltonian

$$H(t) = \frac{1}{2}\omega(t)(S_+ + S_-) \quad (1)$$

in terms of $|+\rangle$ and $|-\rangle$ to describes this transition process of helicity states on the LRH interfaces, where $S_+ = |+\rangle\langle-|$, $S_- = |-\rangle\langle+|$, $S_3 = \frac{1}{2}(|+\rangle\langle+| - |-\rangle\langle-|)$ satisfying the following SU(2) Lie algebraic commuting relations $[S_+, S_-] = 2S_3$ and $[S_3, S_{\pm}] = \pm S_{\pm}$. The time-dependent frequency parameter $\omega(t)$ may be taken to be $\omega(t) = \zeta \frac{d}{dt}p(t)$, where $p(t) = (-)^m$ with $m = [\frac{ct}{nb}]$, and ζ is the coupling coefficient, which is determined by the physical mechanism of interaction between light field and media. Since $p(t)$ is a periodical function, by using the analytical continuation procedure, it can be rewritten as the following combinations of analytical functions

$$p(t) = \sum_{k=1}^{\infty} \frac{2}{k\pi} [1 - (-)^k] \sin\left(\frac{k\pi c}{nb}t\right). \quad (2)$$

In what follows, we solve the time-dependent Schrödinger equation (in the unit $\hbar = 1$)

$$H(t)|\Psi_{\sigma}(t)\rangle = i\frac{\partial}{\partial t}|\Psi_{\sigma}(t)\rangle \quad (3)$$

governing the propagation of light in the LRH- periodical fiber. According to the Lewis-Riesenfeld invariant theory, the exact particular solution $|\Psi_{\sigma}(t)\rangle$ of the time-dependent Schrödinger equation (3) is different from the eigenstate of the invariant $I(t)$ only by a time-dependent c - number factor $\exp[\frac{1}{i}\varphi_{\sigma}(t)]$, where

$$\varphi_{\sigma}(t) = \int_0^t \langle\Phi_{\sigma}(t')|[H(t') - i\frac{\partial}{\partial t'}]|\Phi_{\sigma}(t')\rangle dt' \quad (4)$$

with $|\Phi_{\sigma}(t)\rangle$ being the eigenstate of the invariant $I(t)$ (corresponding to the particular eigenvalue σ) and satisfying the eigenvalue equation $I(t)|\Phi_{\sigma}(t)\rangle = \sigma|\Phi_{\sigma}(t)\rangle$, where the eigenvalue σ of the invariant $I(t)$ is *time-independent*. Thus we have

$$|\Psi_{\sigma}(t)\rangle = \exp\left[\frac{1}{i}\varphi_{\sigma}(t)\right]|\Phi_{\sigma}(t)\rangle. \quad (5)$$

So, in order to obtain $|\Psi_{\sigma}(t)\rangle$, we should first obtain the eigenstate $|\Phi_{\sigma}(t)\rangle$ of the invariant $I(t)$. In accordance with the Lewis-Riesenfeld theory, the invariant $I(t)$ is a conserved operator (*i. e.*, it possesses *time-independent* eigenvalues) and agrees with the following Liouville-Von Neumann equation

$$\frac{\partial I(t)}{\partial t} + \frac{1}{i}[I(t), H(t)] = 0. \quad (6)$$

It follows from Eq. (6) that the invariant $I(t)$ may also be constructed in terms of $S_{\pm}$ and S_3 , *i. e.*,

$$I(t) = 2 \left\{ \frac{1}{2} \sin \theta(t) \exp[-i\phi(t)] S_+ + \frac{1}{2} \sin \theta(t) \exp[i\phi(t)] S_- + \cos \theta(t) S_3 \right\}. \quad (7)$$

Inserting Eq. (1) and (7) into Eq. (6), one can arrive at a set of auxiliary equations

$$\begin{aligned} \exp[-i\phi] \left(\dot{\theta} \cos \theta - i\dot{\phi} \sin \theta \right) - i\omega \cos \theta &= 0, \\ \dot{\theta} + \omega \sin \phi &= 0, \end{aligned} \quad (8)$$

which are used to determine the time-dependent parameters, $\theta(t)$ and $\phi(t)$, of the invariant $I(t)$.

It should be noted that we cannot easily solve the eigenvalue equation $I(t) |\Phi_{\sigma}(t)\rangle = \sigma |\Phi_{\sigma}(t)\rangle$, for the time-dependent parameters $\theta(t)$ and $\phi(t)$ are involved in the invariant (7). If, however, we could find (or construct) a unitary transformation operator $V(t)$ to make $V^{\dagger}(t)I(t)V(t)$ be *time-independent*, then the eigenvalue equation problem of $I(t)$ is therefore easily resolved. According to our experience for utilizing the invariant-related unitary transformation formulation [22], we suggest a following unitary transformation operator

$$V(t) = \exp[\beta(t) S_+ - \beta^*(t) S_-], \quad (9)$$

where $\beta(t)$ and $\beta^*(t)$ will be determined by calculating $I_V = V^{\dagger}(t)I(t)V(t)$ in what follows.

Calculation of $I_V = V^{\dagger}(t)I(t)V(t)$ yields

$$I_V = V^{\dagger}(t)I(t)V(t) = 2S_3, \quad (10)$$

if β and β^* are chosen to be $\beta(t) = -\frac{\theta(t)}{2} \exp[-i\phi(t)]$, $\beta^*(t) = -\frac{\theta(t)}{2} \exp[i\phi(t)]$. This, therefore, means that we can change the time-dependent $I(t)$ into a time-independent I_V , and the result is $I_V = 2S_3$. Thus, the eigenvalue equation of I_V is $I_V |\sigma\rangle = \sigma |\sigma\rangle$ with $\sigma = \pm 1$, and consequently the eigenvalue equation of $I(t)$ is $I(t)V(t)|\sigma\rangle = \sigma V(t)|\sigma\rangle$. So, we obtain the eigenstate $|\Phi_{\sigma}(t)\rangle$ of $I(t)$, *i. e.*, $|\Phi_{\sigma}(t)\rangle = V(t)|\sigma\rangle$.

Correspondingly, $H(t)$ is transformed into

$$H_V(t) = V^{\dagger}(t) \left[H(t) - i \frac{\partial}{\partial t} \right] V(t) \quad (11)$$

and the time-dependent Schrödinger equation (3) is changed into

$$H_V(t) |\Psi_{\sigma}(t)\rangle_V = i \frac{\partial}{\partial t} |\Psi_{\sigma}(t)\rangle_V \quad (12)$$

under the unitary transformation $V(t)$, where $|\Psi_{\sigma}(t)\rangle_V = V^{\dagger}(t) |\Psi_{\sigma}(t)\rangle$.

Further analysis shows that the exact particular solution $|\Psi_{\sigma}(t)\rangle_V$ of the time-dependent Schrödinger equation (12) is different from the eigenstate $|\sigma\rangle$ of the *time-independent* invariant I_V only by a time-dependent *c*-number factor $\exp\left[\frac{1}{i}\varphi_{\sigma}(t)\right]$ [15], which is now rewritten as $\exp\left\{\int_0^t \langle \sigma | H_V(t') | \sigma \rangle dt'\right\}$.

By using the auxiliary equations (8), it is verified that $H_V(t)$ depends only on the operator S_3 , *i. e.*,

$$H_V(t) = \left\{ \omega(t) \sin \theta(t) \cos \phi(t) + \dot{\phi}(t) [1 - \cos \theta(t)] \right\} S_3 \quad (13)$$

and the time-dependent *c*-number factor $\exp\left[\frac{1}{i}\varphi_{\sigma}(t)\right]$ is therefore $\exp\left\{\frac{1}{i}\left[\varphi_{\sigma}^{(d)}(t) + \varphi_{\sigma}^{(g)}(t)\right]\right\}$, where the dynamical phase is

$$\varphi_{\sigma}^{(d)}(t) = \sigma \int_0^t \omega(t') \sin \theta(t') \cos \phi(t') dt' \quad (14)$$

and the geometric phase is

$$\varphi_{\sigma}^{(g)}(t) = \sigma \int_0^t \dot{\phi}(t') [1 - \cos \theta(t')] dt'. \quad (15)$$

Hence the particular exact solution of the time-dependent Schrödinger equation (3) corresponding to the particular eigenvalue, σ , of the invariant $I(t)$ is of the form

$$|\Psi_{\sigma}(t)\rangle = \exp \left\{ \frac{1}{i} \left[\varphi_{\sigma}^{(d)}(t) + \varphi_{\sigma}^{(g)}(t) \right] \right\} V(t) |\sigma\rangle. \quad (16)$$

It follows from the obtained expression (15) for geometric phase of photons that, if the frequency parameter ω is small (*i. e.*, the adiabatic quantum process) and then according to the auxiliary equations (8), $\theta \simeq 0$, the Berry phase (adiabatic geometric phase) in a cycle (*i. e.*, one round trip, $T \simeq \frac{2\pi}{\omega}$) of parameter space of invariant $I(t)$ is

$$\varphi_{\sigma}^{(g)}(T) = 2\pi\sigma(1 - \cos \theta), \quad (17)$$

where $2\pi(1 - \cos \theta)$ is a solid angle over the parameter space of the invariant $I(t)$, which means the geometric phase is related only to the geometric nature of the pathway along which quantum systems evolve. Expression (17) is analogous to the magnetic flux produced by a monopole of strength σ existing at the origin of the parameter space. This, therefore, implies that geometric phases differ from dynamical phases and involve the global and topological properties of the time evolution of quantum systems.

It is worthwhile to point out that the geometric phase of photons due to helicity inversion presented here is of quantum level. However, whether the Chiao-Wu geometric phase due to the spatial geometric shape of fiber is of quantum level or not is not apparent (see, for example, the arguments between Haldane and Chiao, Tomita about this problem [24,25]), since the expression for the Chiao-Wu geometric phase can be derived by using both the classical Maxwell's electromagnetic theory, differential geometry and quantum mechanics [13,14,23–25]. However, the geometric phase in this Letter can be considered only by Berry's adiabatic quantum theory and Lewis-Riesenfeld invariant theory, namely, the classical electrodynamics cannot predict this geometric phase. Although many investigators have taken into account the boundary condition problem and anomalous refraction in left-handed media by using the classical Maxwell's theory [4,6], less attention is paid to this geometric phase due to helicity inversion. It is believed that this geometric phase originates at the quantum level, but survives the correspondence-principle limit into the classical level. So, We emphasize that it is essential to take into consideration this geometric phase in investigating the anomalous refraction on the LRH interfaces. In addition, in the Chiao-Wu case, there exists the vacuum geometric phase at pure quantum level, which arises from the vacuum quantum fluctuations of photons field [18]. We think such quantum vacuum geometric phase deserves consideration also in the present case.

It is well known that geometric phases arise only in time-dependent quantum systems. In the present problem, the transitions between helicity states on the LRH interfaces, which is a time-dependent process, results in the geometric phase of photons. This may be viewed from two aspects: (i) it is apparently seen in Eq. (8) that if the frequency parameter ω in the Hamiltonian (1) vanishes, then $\dot{\phi} = 0$ and the geometric phase (15) is therefore vanishing; (ii) it follows from (2) that the frequency coefficient $\omega(t)$ of Hamiltonian (1) is

$$\omega(t) = \zeta \frac{d}{dt} p(t) = \frac{2c}{nb} \zeta \sum_{k=1}^{\infty} [1 - (-)^k] \cos \left(\frac{k\pi c}{nb} t \right). \quad (18)$$

Since $|\cos(\frac{k\pi c}{nb} t)| \leq 1$, the frequency coefficient, the transition rates between helicity states and the consequent time-dependent phase ($\varphi_{\sigma}^{(g)}(t) + \varphi_{\sigma}^{(d)}(t)$) greatly decrease correspondingly as the periodical optical path nb greatly increases. Thus we can conclude that the interaction of light fields with media on the LRH interfaces gives rise to this topological quantum phase.

In addition to obtaining the expression (15) for geometric phase, we obtain the wavefunction (16) of photons in the LRH- optical fiber by solving the time-dependent Schrödinger equation (3) based on the Lewis-Riesenfeld invariant theory [15] and the invariant-related unitary transformation formulation [16,17]. We believe that this would enable us to consider the propagation of light fields inside the optical fiber in more detail.

Since we have constructed an effective Hamiltonian (1) to describe the time evolution of helicity states of photons, it should be noted that the method presented here is a phenomenological description of propagation of lightwave in the LRH-periodical fiber. This phenomenological description is based on the assumption that the direction of wave vector $\mathbf{k}$ becomes opposite nearly instantaneously on the LRH interfaces. This assumption holds if the periodical length b is much larger than the wavelength of lightwave in the fiber.

To close this Letter, we conclude with some remarks on the potential significance of the subject in this Letter:

- (i) The obtained geometric phase itself is physically interesting. Moreover, it is necessary to consider this geometric phase in discussing the anomalous refraction and wave propagation in left-handed media (on the LRH interfaces).
- (ii) Helicity inversion of photons, which is in exact analogy with the transition operation from 0 to 1 in digital circuit, can be caused due to the electromagnetic interactions on LRH interfaces, and the time evolution of helicity states is governed by (16). We think these processes may possess some potential applications in information technology and therefore deserve further investigation.
- (iii) It is of physical interest to consider the quantum effects such as propagation of photons field, polarization of photon states (time evolution of photon wavefunction) and spontaneous emission decay rate of atoms [2] in left-handed media. In this Letter, an illustrative example of quantum effects of photons field resulting from helicity inversion caused by the LRH interfaces is presented. We hope the present consideration in this paper would open up new opportunities for investigating more quantum mechanical properties, phenomena and effects in left-handed media.

Acknowledgements I thank X. C. Gao and S. L. He for useful discussions. This project is supported by the National Natural Science Foundation of China under the project No. 90101024.

-
- [1] Smith D R, Padilla W J, Vier D C *et al.* 2000 *Phys. Rev. Lett.* **84** 4184
 - [2] Klimov V V 2002 *Opt. Comm.* **211** 183
 - [3] Pendry J B, Holden A J, Robbins D J and Stewart W J 1999 *IEEE Trans. Microwave Theory Tech.* **47** 2075
 - [4] Shelby R A, Smith D R and Schultz S 2001 *Science* **292** 77
 - [5] Ziolkowski R W 2001 *Phys. Rev. E* **64** 056625
 - [6] Veselago V G 1968 *Sov. Phys. Usp.* **10** 509
 - [7] Pendry J B, Holden A J, Stewart W J and Youngs I 1996 *Phys. Rev. Lett.* **76** 4773
 - [8] Pendry J B, Holden A J, Robbins D J and Stewart W J 1998 *J. Phys. Condens. Matter* **10** 4785
 - [9] Maslovski S I, Tretyakov S A and Belov P A 2001 *Inc. Microwave Opt. Tech. Lett.* **35** 47
 - [10] Berry M V 1984 *Proc. Roy. Soc. London Ser. A* **392** 45
 - [11] Chiao R Y and Wu Y S 1986 *Phys. Rev. Lett.* **57** 933
 - [12] Tomita A and Chiao R Y 1986 *Phys. Rev. Lett.* **57** 937
 - [13] Kwiat P G and Chiao R Y 1991 *Phys. Rev. Lett.* **66** 588
 - [14] Robinson A L 1986 *Science* **234** 424
 - [15] Lewis H R and Riesenfeld W B 1969 *J. Math. Phys.* **10** 1458
 - [16] Gao X C, Xu J B and Qian T Z 1991 *Phys. Rev. A* **44** 7016
 - [17] Gao X C, Fu J, Li X H and Gao J 1998 *Phys. Rev. A* **57** 753
 - [18] Shen J Q and Ma L H 2003 *Phys. Lett. A* **308** 355
 - [19] Shen J Q, Zhu H Y and Shi S L 2002 *Acta Phys. Sin.* **51** 536
 - [20] Anandan J 1977 *Phys. Rev. D* **15** 1448
 - [21] Dresden M and Yang C N 1979 *Phys. Rev. D* **20** 1846
 - [22] Shen J Q, Zhu H Y and Mao H 2002 *J. Phys. Soc. Jpn.* **71** 1440
 - [23] Haldane F D M 1986 *Opt. Lett.* **11** 730
 - [24] Haldane F D M 1987 *Phys. Rev. Lett.* **15** 1788
 - [25] Chiao R Y and Tomita A 1987 *Phys. Rev. Lett.* **15** 1789